Pronil 10mg/ml (1%)

DESCRIPTION AND COMPOSITION:

Acepromazine is a neuroleptic belonging to the phenothiazines with a sedative action. The effect is set in within 5 - 10 minutes after intravenous injection and within 15 - 20 minutes after intramuscular injection and is maintained for 6 to 12 hours.

Product type: Injection

Content: Acepromazine 10mg / ml

TARGET SPECIES:

Swine.

INDICATIONS:

All situations where sedation is needed, for instance during transport, with several forms of stress, small operations and as a pre anaesthetic with larger operations.

ROUTE OF ADMINISTRATION:

Intramuscular or intravenous

WITHDRAWAL PERIODS:

Horses: Meat and offal: 5 days. Milk: 5 days.

DOSAGE:

For intramuscular or intravenous administration.
Calves, cattle, horses and pigs: - Light sedation and premedication with general anaesthesia: 0.5 ml per 100 kg body weight intravenously or 1 ml per 100 kg body weight intramuscularly. - Strong sedation: 1 ml per 100 kg body weight intravenously or 2 ml per 100 kg body weight intramuscularly. Dogs and cats: - Light sedation and premedication with general anaesthesia: 0.25 ml per 10 kg body weight intravenously or intramuscularly. - Strong sedation: 0.5 ml per 10 kg body weight intravenously or intramuscularly.



SHELF LIFE:	PACK SIZE:
3 years	100ml

INJECTABLES

Sedatives & Anaesthetics

Manufacturer:
RANGE PHARMA SDN. BHD.
No.1, Jalan TSB 11, Taman Industri Sg. Buloh,
47000 Sg. Buloh, Selangor, Malaysia.

Tel: (603) 6156 8708
Fax: (603) 6156 8707
Email: info@rangepharma.com
Website: www.rangepharma.com