Quinosol

DESCRIPTION AND COMPOSITION:

Enrofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella and Mycoplasma spp.

Product type: Water Soluble Liquid

Content: Enrofloxacin 200mg / ml

TARGET SPECIES:

Poultry.

INDICATIONS:

For use in turkeys, broilers, and pullets being reared as layers for diseases of the respiratory and alimentary tract of bacterial or mycoplasmal origin such as pasteurellosis, mycoplasmosis, colibacillosis, colisepticaemia and Salmonellosis.

ROUTE OF ADMINISTRATION:

Oral, via drinking water

WITHDRAWAL PERIODS:

Chickens and turkeys: 8 days. Do not use in laying birds producing eggs for human consumption.

DOSAGE:

10mg per kg body weight daily. Total number of birds x Average body weight in kg x 0.05 = Total volume of Quinosol 20% per day. For the treatment of Salmonellosis and Pasteurellosis: 5 to 10 days. For other conditions: 3 to 5 days.

Side effects: Administration to young animals during growth can cause cartilage lesions in joints. Hypersensitivity reactions.



SHELF LIFE:	PACK SIZE:
3 years	500ml, 1L

WATER SOLUBLE LIQUIDS

Antibiotics

Manufacturer:
RANGE PHARMA SDN. BHD.
No.1, Jalan TSB 11, Taman Industri Sg. Buloh,
47000 Sg. Buloh, Selangor, Malaysia.

Tel: (603) 6156 8708
Fax: (603) 6156 8707
Email: info@rangepharma.com
Website: www.rangepharma.com