



Our anticoccidials as a solution.



RANGE PHARMA SDN. BHD. (Co. No. 144401U)
No.1 Jalan TSB 11, Taman Industri Sungai Buloh,
47000 Sungai Buloh, Selangor Darul Ehsan, Malaysia.

Email: info@rangepharma.com
Tel : (603) 6156 8708
Fax : (603) 6156 8707

www.rangepharma.com



AMPROFEED 250G/KG POWDER

Composition:

Each kg contains 250g amprolium as its active ingredient.

Indication:

As an aid in the prevention of coccidiosis in turkeys and chickens (broilers and layer replacement hens).

Dosage:

Broilers: 500g - 1kg Amprofeed per tonne feed.

Replacement hens:

0-5 weeks old: 500g Amprofeed per tonne feed

5-9 weeks old: 320-500g Amprofeed per tonne feed

9-14 weeks old: 320g Amprofeed per tonne feed

Withdrawal Period:

0 days.

Remarks:

Do not use amprolium medicated feed for treatment of outbreaks of coccidiosis.



Pack Size:
25kg

DECOFEED 60G/KG POWDER

Composition:

Each kg contains 60g decoquinate as its active ingredient.

Indication:

Chickens: (broilers): For the prevention of coccidiosis caused by Eimeria tenella, E. necatrix, E. mivati, E. acervulina, E. maxima and E. brunetti.

Dosage:

Chickens (broiler): 27.2g per ton of feed.

Withdrawal Period:

0 days.

Remarks:

Bentonite should not be used in decoquinate feeds.



Pack Size:
25kg

DOTFEED 250G/KG POWDER

Composition:

Each kg contains 250g of dinitro-o-toluidamide as its active ingredient.

Indication:

Turkeys (meat) and chickens (broilers): For the prevention and control of coccidiosis.

Chickens (replacement - growers, starters): Development of active immunity to coccidiosis for chickens under light to moderate exposure. Development of active immunity to coccidiosis for chickens under severe exposure.

Dosage:

Chickens (Broilers): 500g of Dotfeed per tonne feed.

Chickens (replacements - growers) - Development of active immunity to coccidiosis with chickens with severe exposure: 500g of Dotfeed per tonne feed.

Withdrawal Period:

0 days.



Pack Size:
25kg

DURACIN 10G/KG POWDER

Composition:

Each kg contains 10g of Maduramicin Ammonium.

Indication:

Chickens: (broilers): For the prevention of coccidiosis caused by Eimeria tenella, E. necatrix, E. mivati, E. acervulina, E. maxima and E. brunetti.

Dosage:

Chickens (broiler): 500g per tonne of feed.

Withdrawal Period:

5 days.



Pack Size:
25kg

NICOXIN 25% POWDER

Composition:

Each kg contains 250g of nicarbazin as its active ingredient.

Indication:

Chickens: (broilers): For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. mivati*, *E. acervulina*, *E. maxima* and *E. brunetti*.

Dosage:

Poultry: 500g of Nicoxin per tonne feed.

Withdrawal Period:

5 days.



Pack Size:
25kg

ROBENXIN 66G/KG POWDER

Composition:

Each kg contains 66g of Robenidine Hydrochloride.

Indication:

Chickens: (broilers): For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. mivati*, *E. acervulina*, *E. maxima* and *E. brunetti*.

Dosage:

500g of Robenxin 66g/kg Powder per tonne feed.

Withdrawal Period:

Chickens (Broilers): 5 days.

Eggs: Do not use in birds which are producing or may in the future produce eggs or egg products for human consumption



Pack Size:
25kg

ZURICOX 0.5% POWDER

Composition:

Each kg contains 5g diclazuril as its active ingredient.

Indication:

As an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mitis*, *E. necatrix* and *E. tenella* in broiler chickens.

Dosage:

Broiler chickens: 200g of Zuricox 0.5% Powder per tonne of feed (equivalent to 1 part per million of diclazuril) continuously.

Withdrawal Period:

Chickens and turkeys:
0 days.

Not for use in animals from which eggs are produced for human consumption.



Pack Size:
25kg