



Our anticoccidials as a solution.



RANGE PHARMA SDN. BHD. (Co. No. 144401U)
No.1 Jalan TSB 11, Taman Industri Sungai Buloh,
47000 Sungai Buloh, Selangor Darul Ehsan, Malaysia.

Email: info@rangepharma.com
Tel : (603) 6156 8708
Fax : (603) 6156 8707

www.rangepharma.com



AMPROFEED 250G/KG POWDER

Composition:

Each kg contains 250g amprolium as its active ingredient.

Indication:

Turkeys and chickens (broilers and layer replacement hens):
As an aid in the prevention of coccidiosis.

Dosage:

Turkeys: 500g of Amprofeed per tonne of feed.
Broilers: 500g to 1kg of Amprofeed per tonne of feed.
Replacement hens:
0-5 weeks old: 500g of Amprofeed per tonne of feed
5-9 weeks old: 320g to 500g of Amprofeed per tonne of feed
9-14 weeks old: 320g of Amprofeed per tonne of feed

Withdrawal Period:

Turkeys and chickens: 0 days.
Do not feed to laying birds in production.

Remarks:

Do not use amprolium medicated feed for treatment of outbreaks of coccidiosis.



Pack Size:
25kg

CLOPIFEED 250G/KG POWDER

Composition:

Each kg contains 250g of clodol as its active ingredient.

Indication:

Chickens (broilers and layer replacements): Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*.

Dosage:

Chickens (broilers and layer replacements): 500g of Clopifeed 250g/kg Powder per tonne of feed.

If intended for use as caged layers: 1kg of Clopifeed 250g/kg Powder per tonne of feed. Feed up to 16 weeks of age continuously.

Withdrawal Period:

Chicken (broilers): 5 days before slaughter if given at the level of 0.025 percent in feed or reduce level to 0.0125 percent.



Pack Size:
25kg

DECOFEED 60G/KG POWDER

Composition:

Each kg contains 60g decoquinate as its active ingredient.

Indication:

Chickens (broilers): For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. mivati*, *E. acervulina*, *E. maxima* and *E. brunetti*.

Dosage:

Chickens (broiler): 500g of Decofeed 60g/kg Powder per tonne of feed per day.

Withdrawal Period:

Do not feed to laying hens producing eggs for human consumption.

Remarks:

Bentonite should not be used in decoquinate feeds.



Pack Size:
25kg

DOTFEED 250G/KG POWDER

Composition:

Each kg contains 250g of dinitolmide as its active ingredient.

Indication:

Chickens (broilers): For the prevention of coccidiosis.

Chickens (replacement - growers, starters): Development of active immunity to coccidiosis for chickens under light to moderate exposure. Development of active immunity to coccidiosis for chickens under severe exposure.

Dosage:

Chickens (broilers): 500g of Dotfeed 250g/kg Powder per tonne of feed.

Chickens (replacements - growers): Development of active immunity to coccidiosis with chickens with severe exposure: 500g of Dotfeed 250g/kg Powder per tonne of feed.

Withdrawal Period:

0 days.



Pack Size:
25kg

DURACIN 10G/KG POWDER

Composition:

Each kg contains 10g of maduramicin ammonium as its active ingredient.

Indication:

Chickens (broilers): For the prevention of coccidiosis in broiler chickens caused by *Eimeria acervulina*, *E. tenella*, *E. brunetti*, *E. maxima*, *E. necatrix* and *E. mivati*.

Dosage:

Chickens (broiler): 450g to 550g of Duracin 10g/kg Powder per tonne of feed per day. For 5 consecutive days orally in feed.

Withdrawal Period:

5 days.



Pack Size:
25kg

NICOXIN 25% POWDER

Composition:

Each kg contains 250g of nicarbazin as its active ingredient.

Indication:

Chickens (broilers): For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. tenella*, *E. brunetti*, *E. maxima*, *E. necatrix* and *E. mivati*.

Dosage:

500g of Nicoxin 25% Powder per tonne of feed.

Withdrawal Period:

5 days.



Pack Size:
25kg

ROBENXIN 66G/KG POWDER

Composition:

Each kg contains 66g of robenidine hydrochloride as its active ingredient.

Indication:

Chickens (broilers): For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. tenella*, *E. brunetti*, *E. maxima*, *E. necatrix* and *E. mivati*.

Dosage:

500g of Robenxin 66g/kg Powder per tonne of feed.

Withdrawal Period:

Chickens (broilers): 5 days.

Do not feed to laying hens producing eggs for human consumption.



Pack Size:
25kg

ZURICOX 0.5% POWDER

Composition:

Each kg contains 5g diclazuril as its active ingredient.

Indication:

Chickens (broilers): For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. mivati*, *E. acervulina*, *E. maxima* and *E. brunetti*.

Turkeys: For the prevention of coccidiosis caused by *Eimeria adenoeides*, *E. meleagrimitis* and *E. gallopavonis*.

Dosage:

Broiler chickens: 200g of Zuricox 0.5% Powder per tonne of feed.

Growing turkeys: 200g of Zuricox 0.5% Powder per tonne of feed with bacitracin methylene disalicylate 4g to 50g per tonne of feed continuously as the sole ration.

Withdrawal Period:

Chickens (broilers) and turkeys: 0 days.

Do not feed to animals producing eggs for human consumption.



Pack Size:
25kg