

Product Catalogue

We're All About Animal Health










Table of Contents

Injections		02 - 16
Water Soluble Liquid		17 - 23
Water Soluble Powder		24 - 34
Oral Pumps		35 - 36
Feed Additives		37 - 47
Disinfectants		48 - 51

INJECTIONS

ADE Fort

Indications

For the treatment and prevention of vitamin A, D and E deficiencies in cattle, sheep and pigs.

Dosages

By intramuscular injection only.

Cows : 5ml at least twice per year and 30 days prior to calving.

Calves : 1ml during the first week after birth.

Sheep : 1ml to 2ml: At least twice per year.

Lambs : 0.5ml: As soon as possible after birth.

Pigs : 2ml to 3ml;

Sows : One injection upon placement in farrowing house and a second injection during the first week after farrowing.

Boars : One injection 1 to 2 weeks prior to breeding season.

Piglets : 0.25ml to 0.5ml during the first week after birth.

API: Vitamin A 500,000 IU/ml, Vitamin D3 50,000 IU/ml, Vitamin E 50mg/ml



Pack size: 100ml

Azanil 40mg/ml

Indications

A neuroleptic sedative for pigs to be used for the treatment of:

- i) Aggression
 - Prevention of fighting
 - Treatment of aggression in sows
- ii) Stress, including transport-related stress.
- iii) Obstetric conditions e.g., cessation of parturition due to excitation, as an obstetric aid in manual delivery, inversion of the vagina, prolapse of the uterus, pathological straining.
- iv) Premedication in local and general anaesthesia.

Dosages

To be given strictly by intramuscular injection, behind the ear.

General Transportation: 0.5ml to 1ml / 100kg body weight

Transport of boars: 0.5ml / 20kg body weight

Prevention of fighting, treatment of aggression in sows, obstetrics: 1ml / 20kg body weight

Restlessness, anxiety, nervousness, excitation, premedication in local and general anaesthesia: 0.5ml to 1ml / 20kg body weight

API: Azaperone 40mg/ml



Pack size: 100ml

Balamin 0.1%

Indications

Prevention and treatment of vitamin B12 deficiency in cattle and pigs.

Dosages

Intravenous, intramuscular and subcutaneous injection.
Cattle and pigs: 1ml to 2ml / 100kg body weight. These doses can be restarted each daily for 8 days.

API: Cyanocobalamin (Vitamin B12) 1mg/ml



Pack size: 100ml

Clamoxon SP

Indications

Treatment of respiratory, gastrointestinal, biliary and urinary tract infections. It is also indicated for the treatment of infection of the uterus and navel as well as mastitis and secondary infections.

Dosages

Cattle and pigs: 15mg of amoxicillin and gentamicin 4mg / kg body weight equivalent to 1ml of Clamoxon SP / 10kg body weight for 3 days via intramuscular injection.

API: Amoxicillin Trihydrate 150mg/ml, Gentamicin 40mg/ml



Pack size: 100ml

Dexavet 0.5%

Indications

Anti-inflammatory and anti-allergic agent in horses, cattle and pigs.

Dosages

Horses, cattle and pigs: 0.6ml / 50kg body weight injected intramuscularly. Repeat once at 24 to 48 hours intervals, if necessary.

API: Dexamethasone 5mg/ml



Pack size: 100ml

Fluniflor

Indications

Treatment of bovine respiratory disease complex (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni* and BRD associated pyrexia.

Dosages

A single subcutaneous injection of Fluniflor at 2ml / 15kg body weight, providing 40mg / kg florfenicol and 2.2mg / kg flunixin.

The injection should only be given in the neck.

API: Florfenicol 300mg/ml, Flunixin 16.5mg/ml



Pack size: 100ml

Flunivet 5%

Indications

Cattle: Reduction of acute inflammation associated with respiratory disease. Adjunctive therapy in the treatment of acute mastitis.

Horses: Alleviation of inflammation and pain associated with musculoskeletal disorders. Alleviation of visceral pain associated with colic. Adjunctive therapy in the treatment of endotoxaemia and septic shock.

Pigs: Adjunctive therapy in the treatment of swine respiratory diseases.

Dosages

Cattle: 2ml / 45kg body weight administered intravenously. Repeat as necessary at 24 hour intervals for up to 5 consecutive days.

Horses: By intravenous injection.

For colic: 1ml / 45kg body weight repeated once or twice if colic recurs.

For musculoskeletal disorders: 1ml / 45kg body weight once daily for up to 5 days according to clinical response.

For the treatment of endotoxaemia or septic shock associated with gastric torsion and with other conditions in which the circulation of blood to the gastrointestinal tract is compromised: 0.25mg / kg every 6 to 8 hours.

Pigs: 2ml / 45kg body weight once by intramuscular injection, in the neck, in conjunction with appropriate antimicrobial therapy. The injection volume should be limited to a maximum of 5ml / injection site.

API: Flunixin (as Meglumine) 50mg/ml



Pack size: 100ml

Florovet 400mg/ml

Indications

Cattle: For the treatment of *Pasteurella pneumonia*, *Haemophilus pneumonia*, nephropathy caused by *Pasteurella haemolytica*, *Pasteurella multocida*, *Haemophilus somnus*, *Fusobacterium necrophorum*, and *Bacteroides melanogenicus* susceptible to florfenicol.

Pigs: For the treatment of Pleural pneumonia, *Pasteurella pneumonia*, *Mycoplasma pneumonia*, *Haemophilus pneumonia*, *Salmonella* diarrhoea caused by *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Haemophilus parasuis*, *Salmonella* spp., *Mycoplasma hyopneumoniae* and *Mycoplasma hyorhinis* susceptible to florfenicol.

Dosages

Cattle: 0.05ml of product (equivalent to 20mg florfenicol) / kg body weight intramuscularly into the neck once a day, repeat the same amount after 48 hours.

Pigs: 0.037ml of product (equivalent to 15mg florfenicol) / kg body weight intramuscularly once a day into the neck area, repeat the same amount after 48 hours. Do not exceed 3ml per injection site.

API: Florfenicol 400mg/ml



Pack size: 100ml

Gentacin 10%

Indications

Chickens: Prevention of early mortality involving *E. coli*, *Salmonella thimurium* and *Pseudomonas aeruginosa* susceptible to gentamicin.

Pigs: Treatment of porcine colibacillosis caused by *E. coli* sensitive to gentamicin.

Dosages

Chickens: 1ml of Gentacin 10% diluted to 99ml of sterile saline solution and injected aseptically at 0.2ml / day old chicken subcutaneously in the neck.

Piglets: 2.5ml of Gentacin 10% diluted to 47.5ml of sterile saline solution and injected aseptically at 1ml / piglet intramuscularly.

API: Gentamicin (as Sulphate) 100mg/ml



Pack size: 100ml

Gleptofort 200mg/ml

Indications

For prevention and treatment of iron deficiency anaemia in piglets.

Dosages

Strictly for intramuscular injection.

Piglets: 1ml of the product which is equivalent to 200mg Iron (III) per animal. Inject once between the 1st and the 3rd day of life.

API: Iron (as Gleptoferron) 200mg/ml



Pack size: 100ml

Hemofort 20%

Indications

Prevention and treatment of iron deficiency anaemia in pigs.

Dosages

1ml / piglet by deep intramuscularly injection preferably in the thigh muscle in the three days after birth.

API: Iron (as Dextran) 200mg/ml



Pack size: 100ml

Ironex 10%

Indications

Prevention and treatment of iron deficiency anaemia in pigs.

Dosages

2ml / piglet by deep intramuscular injection preferably in the thigh muscle in the three days after birth.

API: Iron (as Dextran) 100mg/ml



Pack size: 100ml

Ironfort + B12

Indications

Treatment and prevention of iron deficiency anaemia in piglets and calves.

Dosages

For intramuscular route.

Piglets: 100mg to 200mg of iron dextran / piglet, i.e. 1ml to 2ml all at once during one of the first days of the piglet's life, preferably in the muscles of the neck.

Calves: 300mg to 400mg of iron dextran / calf, i.e. 3ml to 4ml all at once during one of the first days of the calf's life, preferably in the muscles of the neck.

Duration of treatment: One injection to be repeated if necessary after 8 days to 1 month.

API: Iron Dextran 100mg/ml, Vitamin B12 0.08mg/ml



Pack size: 100ml

Ivertin 1%

Indications

Treatment of bovine respiratory disease complex (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni* and BRD associated pyrexia.

Dosages

Cattle: 0.02ml / kg body weight by subcutaneous injection under the loose skin in front of or behind the shoulder.

Pigs: 0.03ml / kg body weight by subcutaneous injection into the neck.

Young pigs: Less than 0.03ml / kg body weight.

API: Ivermectin 10mg/ml



Pack size: 100ml

Kanacin 10%

Indications

For the treatment of bacterial infections caused by staphylococcus, Corynebacterium, *E. coli*, Salmonella and Proteus.

Cattle: For treatment of Pneumonia, Bronchitis, bacterial diarrhoea, bacterial arthritis and mastitis.

Pigs: For treatment of Pneumonia, bacterial diarrhoea.

Dosages

For intramuscular administration.

Cattle: 5mg to 10mg kanamycin / kg body weight / day.

Pigs: 10mg to 20mg kanamycin / kg body weight / day.

API: Kanamycin (as Sulphate) 100mg/ml



Pack size: 100ml

Lidonil 20mg/ml

Indications

Horse, cats and dogs: For local/nerve block (regional infiltration) including field block anaesthesia. Superficial anaesthesia of mucous membranes.

Dosages

Horses, cats and dogs: For subcutaneous, intramuscular or perineural injection or for application onto the mucosa. To avoid intravascular administration.

Local/nerve block anaesthesia in horses: 1ml to 10ml of product.

Superficial anaesthesia of mucous membranes: Instil a thin layer topically to site where anaesthesia is required.

The total dose shall not exceed 2mg to 4mg lidocaine HCl / kg body weight (equivalent to 1ml of the product / 5kg to 10kg body weight).

API: Lidocaine HCl 20mg/ml



Pack size: 100ml

Linco Actin

Indications

Calves: For the treatment of respiratory infections, arthritis and omphalitis.

Sheep and goats: For the treatment of respiratory infections and mycoplasma.

Pigs: For the treatment of intestinal adenomatosis (ileitis) caused by *Lawsonia intracellularis*, haemorrhagic enteritis and colibacillosis, mycoplasma and infectious arthritis.

Poultry: For the treatment of mycoplasma associated or not with *E. coli*, airsacculitis caused by *E. coli* and avian cholera caused by *Pasteurella multocida*.

Dogs and cats: For the treatment of respiratory, intestinal, urinary, skin (including wounds and abscesses) and arthritis infections caused by organisms susceptible to lincomycin and/or spectinomycin.

Dosages

Calves: 1ml / 10kg body weight intramuscularly. Two injections at 12 hours intervals on the first day of treatment, then one injection / day for 3 to 5 days.

Sheep and goats: 1ml / 10kg body weight intramuscularly, once daily for 3 to 5 days.

Pigs: 1ml / 10kg body weight intramuscularly, repeat if necessary after 24 hours and for up to 5 days.

Poultry: 1ml / 5kg body weight subcutaneously, once daily for 3 days.

Dogs and cats: 1ml / 5kg body weight intramuscularly. Repeat at 12 and 24 hours intervals for 3 to 7 days depending on clinical response.

API: Lincomycin (as HCl) 50mg/ml, Spectinomycin (as Sulphate) 100mg/ml



Pack size: 100ml

Marboxin 2%

Indications

Treatment of respiratory infections caused by susceptible strains of organisms in cattle and pigs.

Dosages

2mg / kg body weight / day (1ml Marboxin 2% / 10kg) in a single daily injection by subcutaneous or intravenous route in cattle and by intramuscular route in pigs.

Treatment duration is 3 days by subcutaneous route and 3 to 5 days by intravenous route in cattle and 3 to 5 days by intramuscular route in pigs

API: Marbofloxacin 20mg/ml



Pack size: 100ml

Marboxin 10%

Indications

Cattle: Treatment of respiratory infections and for the treatment of acute *E. coli* mastitis.

Pigs: Treatment of Metritis Mastitis Agalactia syndrome (MMA).

Dosages

Administer 1ml / 50kg in a single daily injection by intramuscular, subcutaneous or intravenous routes in cattle and by intramuscular route in pigs. Treatment duration is 3 days in pigs and 3 to 5 days in cattle.

API: Marbofloxacin 100mg/ml



Pack size: 100ml

Moxi Fort 20%

Indications

Effective against a wide range of Gram-positive and Gram-negative bacteria. For treatment of infections of the alimentary tract, respiratory tract, skin and soft tissues, urogenital tract and in prevention of post-operative infection (treat before surgery).

Dosages

Bovine, pigs and sheep: 0.75ml / 10kg body weight. Repeat after 48 hours, if necessary.

API: Amoxicillin (as Trihydrate) 200mg/ml



Pack size: 100ml

Moxi LA 15%

Indications

Effective against a wide range of Gram-positive and Gram-negative bacteria. For treatment of infections of the alimentary tract, respiratory tract, skin and soft tissues, urogenital tract and in prevention of post-operative infection (treat before surgery).

Dosages

Cattle, sheep and pigs: By intramuscular injection only.
Dogs and cats: By subcutaneous or intramuscular injection.
1ml / 10kg body weight. Repeat after 48 hours, if necessary.

API: Amoxicillin (as Trihydrate) 150mg/ml



Pack size: 100ml

Pronil 10mg/ml

Indications

For use in cattle, sheep, swine, goats and horses, as an aid in tranquilising animals:

- Before loading
- To facilitate restraint
- To control hyperexcitability
- As a pre-anaesthetic agent in major and minor surgery

In colts: As an aid in breaking colts.

In bulls and cows: To tranquilise animals for artificial insemination procedures.

For use in dogs and cats as an aid in tranquilising animals:

- To facilitate restraint
- As an aid in the control of hyperexcitability, vomiting, licking and scratching
- As a pre-anaesthetic agent in major and minor surgery
- As an aid in breeding high-strung bitches

Dosages

Sheep and goats:

Sedation: Administer 0.05mg to 0.1mg / kg body weight intramuscularly.

Cattle:

Sedation: 0.01mg to 0.02mg / kg body weight intravenously or 0.03mg to 0.1mg / kg body weight intramuscularly.

Sedation prior to local anaesthesia: 0.1mg / kg body weight intramuscularly.

Swine:

Sedation: Administer 0.01mg to 0.2mg / kg body weight intravenously or intramuscularly.

Dogs:

Sedation: 0.025mg to 0.2mg / kg body weight intravenously to a maximum dose of 3mg intravenously or 0.1mg to 0.25mg / kg body weight intramuscularly or subcutaneously.

Pre-anaesthetic: 0.03mg to 0.2mg / kg body weight intravenously, intramuscularly or subcutaneously, with a maximum dose of 3mg intravenously.

Cats:

Sedation: 0.05mg to 0.11mg / kg body weight intravenously, intramuscularly or subcutaneously, with a maximum dose of 1mg intravenously.

Horses:

Pre-anaesthetic: 0.02mg to 0.05mg / kg body weight intravenously.

Tranquilisation: 0.04mg to 0.1mg / kg body weight intramuscularly.

Dosages may be repeated in 10 hours. As a preloading dose for cattle, sheep, goats, horses and swine, administer 25mg / 454kg intravenously or 75mg / 454kg intramuscularly one hour before loading.

API: Acepromazine Maleate 10mg/ml



Pack size: 100ml

Quinofort 100mg/ml

Indications

Cattle: For the treatment of respiratory tract infections caused by enrofloxacin-sensitive *Histophilus somni*, *Mannheimia haemolytica*, *Pasteurella multocida* and *Mycoplasma* spp. For the treatment of mastitis caused by enrofloxacin-sensitive *E. coli*.

Pigs: For the treatment of bacterial bronchopneumonia caused by enrofloxacin-sensitive *Actinobacillus pleuropneumoniae*, *Pasteurella multocida* and complicated by *Haemophilus parasuis* as a secondary pathogen in pigs.

Dosages

Cattle:

For respiratory diseases: 7.5ml of the product / 100kg body weight / day for a single treatment by subcutaneous injection. In case of serious or chronic respiratory disease a second injection may be required after 48 hours.

For the treatment of colimastitis: 5ml of the product / 100kg body weight / day by intravenous injection. The treatment of colimastitis should be exclusively by intravenous application on 2 to 3 consecutive days.

Pigs:

For respiratory tract infections: 0.75ml of the product / 10kg body weight / day for a single treatment by intramuscular injection into the neck muscles behind the ear. Do not administer more than 7.5ml per injection site. In cases of serious or chronic respiratory disease, a second injection may be required after 48 hours.

API: Enrofloxacin 100mg/ml



Pack size: 100ml

Quinovet 10%

Indications

Cattle: Treatment of diseases of the respiratory diseases and alimentary tract of bacterial or mycoplasmal origin and secondary bacterial infections subsequent to viral conditions (e.g viral pneumonia). Treatment of local signs (inflammation, milk quality and yield) associated with peracute/acute mastitis in lactating dairy cattle caused by *E. Coli*.

Pigs: Treatment of diseases of the respiratory and alimentary tract of bacterial or mycoplasmal origin. Multifactorial diseases such as atrophic rhinitis and enzootic pneumonia.

Dosages

Cattle and pigs: 2.5ml / 100kg body weight daily for 3 days. This rate may be doubled (5ml / 100kg) for 5 days for salmonellosis and complicated respiratory disease.

Cattle: By subcutaneous injection.

Pigs: By intramuscular injection.

API: Enrofloxacin 100mg/ml



Pack size: 100ml

Seftiovet 5%

Indications

For the treatment of infections associated with bacteria sensitive to ceftiofur.

Cattle: For the treatment of bacterial respiratory disease associated with *P. multocida*, *M. haemolytica* and *H. somi*. For the treatment of acute interdigital necrobacillosis (panaritium, foot rot), associated with *F. necrophorum* and *B. melaninogenicus* (*P. asaccharolytica*).

For treatment of the bacterial component of acute post-partum (puerperal) metritis within 10 days after calving associated with *E. coli*, *A. pyogenes* and *F. necrophorum*: this indication is restricted to cases where treatment with another antimicrobial has failed.

Pigs: For the treatment of bacterial respiratory disease associated with *P. multocida*, *A. pleuropneumoniae* and *S. suis*.

Dosages

Cattle: Subcutaneous use.

Respiratory disease: 1mg ceftiofur (as HCl) / kg / day for 3 to 5 days, i.e. 1ml / 50kg at each injection.

Acute interdigital necrobacillosis: 1mg ceftiofur (as HCl) / kg / day for 3 days, i.e. 1ml / 50kg at each injection.

Acute post-partum metritis within 10 days after calving: 1mg ceftiofur (as HCl) / kg / day for 5 consecutive days, i.e. 1ml / 50kg at each injection.

Pigs: Intramuscular use. 3mg ceftiofur (as HCl) / kg / day for 3 days, i.e. 1ml / 16kg at each injection.

API: Ceftiofur (as HCl) 50mg/ml



Pack size: 100ml

Supracin L.A. 20%

Indications

Cattle: For the treatment of respiratory infections caused by oxytetracycline susceptible organisms such as *Arcanobacterium* (Actinomyces) *pyogenes* and *Haemophilus somnus*.

Pigs: For the treatment of respiratory infections caused by oxytetracycline susceptible organisms such as *Pasteurella multocida*.

Dosages

Cattle not producing milk for human consumption:
20mg oxytetracycline / kg body weight (1ml Supracin L.A. 20% / 10kg body weight), if necessary repeat after 72 hours.

Cattle producing milk for human consumption:
20mg oxytetracycline / kg body weight (1ml Supracin L.A. 20% / 10kg body weight) as a single injection only.

Pigs: 20mg oxytetracycline / kg body weight (1ml Supracin L.A. 20% / 10kg body weight), if necessary repeat after 72 hours.

API: Oxytetracycline (as Dihydrate) 200mg/ml



Pack size: 100ml

Taikocin 30%

Indications

Cattle: Treatment of respiratory tract infections due to *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*.

Pigs: Treatment of acute outbreaks of swine respiratory disease caused by strains of *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

Dosages

By intramuscular injection into the neck.

Cattle: 1ml / 15kg body weight, twice 48 hours apart.

Pigs: 1ml / 20kg body weight, twice 48 hours apart.

API: Florfenicol 300mg/ml



Pack size: 100ml

Tolcid 4%

Indications

Cattle: Adjuvant treatment of pneumonia by improving general conditions and runny nose. Adjuvant treatment of acute mastitis.

Pigs: Adjuvant treatment of Metritis Mastitis Agalactia syndrome.

Dogs: Treatment of inflammation associated with musculoskeletal disorders and reduction of post-operative pain.

Cats: Adjuvant treatment of upper respiratory diseases, in combination with antimicrobial treatment, if necessary.

Dosages

Cattle:

Respiratory diseases: 1ml / 20kg body weight intramuscularly in the neck region. Repeat once after 48 hours.

Acute mastitis: 1ml / 10kg body weight in a single slow intravenous injection. At the first signs of intolerance the injection should be stopped.

Pigs: 1ml / 20kg body weight as a single intramuscular injection in the neck.

Cats and dogs: 1ml / 10kg body weight, in a single injection, repeat if necessary, once after 24 to 48 hours depending on clinical assessment. Or a single injection of 1ml / 10kg body weight, followed by oral treatment with tablets.

Dogs: By intramuscular or subcutaneous route.

Cats: By subcutaneous route.

To reduce post-operative pain, it is advisable to perform a pre-operative injection, as a premedication, one hour before induction of anaesthesia.

API: Tolfenamic Acid 40mg/ml



Pack size: 100ml

Triamcort 0.2%

Indications

For the treatment of inflammation and related disorders. It is also indicated for use in the management and treatment of acute arthritis, allergic and dermatologic disorders.

Dosages

The dose is a single injection of 0.06ml to 0.11ml of Triamcort 0.2% Injection / kg of body weight in inflammatory or allergic disorders and 0.11ml of Triamcort 0.2% Injection / kg of body weight in dermatologic disorders.

API: Triamcinolone Acetonide 2mg/ml



Pack size: 100ml

Trimetovet 24%

Indications

Indicated for the treatment of systemic infections caused by or associated with organisms sensitive to the trimethoprim: sulfadiazine combination. The spectrum of activity includes both Gram-positive and Gram-negative organisms including: Actinobacilli, Actinomycae, *Bordetella* spp., Brucella, Corynebacteria, *Escherichia coli*, *Haemophilus* spp., *Klebsiella* spp., *Pasteurella* spp., Pneumococci, Proteus, *Salmonella* spp., Staphylococci, Streptococci and Vibrio.

Dosages

Cattle and pigs: 1ml of product / 16kg body weight by intramuscular or slow intravenous injection. By intravenous injection when rapid blood levels of active ingredients are required.

Horses: 1ml of product / 16kg body weight by slow intravenous injection.

Dogs and cats: 1ml of product / 8kg body weight, by subcutaneous injection only. The recommended site in dogs is the loose skin at the top of the neck. In severe infections it may be repeated daily until 2 days after symptoms resolve, up to a maximum of 5 days.

API: Sulphadiazine 200mg/ml, Trimethoprim 40mg/ml



Pack size: 100ml

Tylocilin 20%

Indications

For the treatment of diseases in pigs involving organisms sensitive to tylosin, such as swine erysipelas (*Erysipelothrix rhusiopathiae*) and pneumonia (*Mycoplasma hyopneumoniae*).

Dosages

0.5ml Tylocilin 20% / 10kg body weight, equivalent to 10mg of tylosin / kg body weight, by deep intramuscular injection every 12 hours, up to a maximum of 6 injections. Do not inject more than 5ml at a single injection site.

API: Tylosin Base 200mg/ml



Pack size: 100ml

Xylanil 20mg/ml

Indications

Sedation of dogs, cats, horses and cattle.

Dosages

Accurately ascertain the body weight of an animal before treatment with xylazine. Use a syringe with appropriate gradations.

Dogs: 0.5ml to 1.0ml of product / 10kg body weight intramuscularly or subcutaneously. 0.35ml to 0.5ml of product / 10kg body weight intravenously.

Cats: 0.125ml to 0.25ml of product / 5kg body weight intramuscularly or subcutaneously.

Horses: 2.5ml to 5.0ml of product / 100kg body weight intravenously.

Cattle: 0.25ml to 1.0ml of product / 100kg body weight intramuscularly. 0.15ml to 0.5ml of product / 100kg body weight intravenously. The intravenous injection should be given slowly, especially in horses.

API: Xylazine (as HCl) 20mg/ml



Pack size: 100ml

Xylanil 100mg/ml

Indications

Cats, dogs and horses: To produce sedation, as an analgesic, and as a preanaesthetic to local or general anaesthesia.

Dosages

Cats, dogs and horses: 0.5ml of product / 45kg body weight intravenously or to 1.0ml of product / 45kg body weight intramuscularly. body weight intravenously. The intravenous injection should be given slowly, especially in horses.

API: Xylazine (as HCl) 100mg/ml



Pack size: 100ml

WATER SOLUBLE LIQUID

Aminosol

Indications

Aminosol contains high concentration of 18 amino acids and 9 minerals for optimal livestock growth.


Dosages

Poultry:

Critical period: 1L to 500 Gallons (2000L) of drinking water.

Normal period: 1L to 1000 Gallons (4000L) of drinking water.

API: Lysine 10.9g/L, Methionine 2.4g/L, Cystine 0.3g/L, Threonine 5.6g/L, Tryptophan 2g/L, Leucine 10g/L, Isoleucine 6.6g/L, Valine 7g/L, Phenylalanine 5.5g/L, Histidine 2.8g/L, Arginine 2.9g/L, Aspartic Acid 12.6g/L, Serine 5.6g/L, Glutamic Acid 17.6g/L, Glycine 6.3g/L, Alanine 9.1g/L, Tyrosine 4.6g/L, Proline 6.3g/L, Calcium 200mg/L, Phosphorus 3.72g/L, Sodium 684mg/L, Magnesium 436mg/L, Potassium 4.84mg/L, Copper 200mcg/L, Zinc 12.2mg/L, Iron 16mg/L, Manganese 2.5mg/L



Pack size: 1L

Aminovit Plus

Indications

Aminovit Plus contains 8 vitamins, 18 amino acids, 9 minerals, immunostimulating peptides and betaine for optimal livestock growth.

Dosages

Poultry:

Critical period: 1L to 500 Gallons (2000L) of drinking water.

Normal period: 1L to 1000 Gallons (4000L) of drinking water.

API: Vitamin A 2,500,000 IU/L, Vitamin E 3750 IU/L, Vitamin B2 Phosphate 1g/L, Vitamin D3 500,000 IU/L, Vitamin B6 125mg/L, Vitamin B12 5mg/L, Biotin 2.5mg/L, Dexpanthenol 300mg/L, Betaine 400mg/L, Lysine 5.5g/L, Methionine 0.5g/L, Cystine 0.1g/L, Threonine 2.8g/L, Tryptophan 1.0g/L, Leucine 5g/L, Isoleucine 3.3g/L, Valine 3.5g/L, Phenylalanine 2.7g/L, Histidine 1.4g/L, Arginine 1.5g/L, Aspartic Acid 6.3g/L, Serine 2.8g/L, Glutamic Acid 8.8g/L, Glycine 3.1g/L, Alanine 4.6g/L, Tyrosine 2.3g/L, Proline 3.2g/L, Calcium 0.1g/L, Phosphorus 1.9g/L, Sodium 342mg/L, Magnesium 218mg/L, Potassium 2.4mg/L, Copper 0.1mg/L, Zinc 6.1mg/L, Iron 8mg/L, Manganese 1.3mg/L, Peptides 5mg/L



Pack size: 1L

Aviacox 2.5%

Indications

For the treatment of coccidiosis in chickens and turkeys:

Chickens: *E. acervulina*, *E. brunetti*, *E. maxima*, *E. mitis*, *E. necatrix* and *E. tenella*.

Turkeys: *E. adenoides* and *E. meleagrimitis*.

Dosages

Chickens and turkeys: 0.28ml of product / kg body weight / day. Treatment is carried out on 2 consecutive days.

API: Toltrazuril 25mg/ml



Pack size: 500ml, 1L

Coxicin

Indications

Treatment of gastrointestinal infections caused by coccidia in poultry.

Dosages

15ml / 10 litres of drinking water for 2 consecutive days. Stop treatment for 3 days then treat for another 2 days.

API: Sulphaquinoxaline (as Sodium) 30mg/ml, Primethamine 10mg/ml



Pack size: 1L

Flumecin 20%

Indications

Poultry: For the treatment and prevention of respiratory infections, gastrointestinal infections and septicaemia caused by *E. coli*, *Salmonella* spp. and *Pasteurella* spp.

Calves: For the treatment and prevention of gastrointestinal infections caused by *E. coli* and *Salmonella* spp. and respiratory infections due to *Pasteurella multocida* and *Mannheimia haemolytica*.

Dosages

Poultry and calves: 12mg flumequine / kg body weight / day for 5 days orally.

API: Flumequine 200mg/ml



Pack size: 1L

Levasol 7.5%

Indications

Broad spectrum anthelmintic for the treatment and control of gastrointestinal and pulmonary nematode infections in sheep and cattle, effective against mature and developing immature stages of the levamisole-susceptible nematode worm species.

Dosages

Levasol 7.5% should be administered as an oral drench. Dosing should be carried out accurately at a rate of 7.5mg levamisole HCl / kg body weight.

Cattle: 1ml Levasol 7.5% / 10kg body weight

Sheep: 0.5ml Levasol 7.5% / 5kg body weight

API: Levamisole HCl 75mg/ml



Pack size: 1L

Micosol 25%

Indications

Pigs: For the treatment and prevention of respiratory infections associated with *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and *Actinobacillus pleuropneumoniae* when the disease has been diagnosed at the herd level.

Chickens: For the treatment and prevention of respiratory infections in poultry flocks associated with *Mycoplasma gallisepticum* and *Mycoplasma synoviae* when the disease has been diagnosed at the herd level.

Turkeys: For the treatment and prevention of respiratory infections in turkey flocks associated with *Mycoplasma gallisepticum* and *Mycoplasma synoviae* when the disease has been diagnosed at the herd level.

Calves: For the treatment and prevention of respiratory infections associated with *Mannheimia haemolytica*, *P. multocida*, *Mycoplasma bovis* and *M. dispar* when the disease has been diagnosed at the herd level. *P. multocida*, *Mycoplasma bovis* and *M. dispar* when the disease has been diagnosed at the herd level.

Dosages

Pigs: 6ml to 8ml of product for 100kg body weight corresponding to 80ml of product / 100 litres of drinking water for 5 days.

Chickens: 6ml to 8ml of product for 100kg body weight corresponding to 30ml of product / 100 litres of drinking water for 3 days.

Turkeys: 4ml to 11ml of product for 100kg body weight corresponding to 30ml of product / 100 litres of drinking water for 3 days.

Calves: 1ml of product for 20kg body weight two times / day for 3 to 5 days.

API: Tilmicosin 250mg/ml



Pack size: 500ml, 1L

Mucolin 1%

Indications

Aids in the treatment of respiratory diseases in cattle, swine and poultry where mucus is a complicating factor.

Dosages

Cattle: 5ml / 100kg body weight once daily for 5 consecutive days.

Pigs: 2ml to 5ml / 100kg body weight once daily for 5 consecutive days.

Poultry: 1ml / litre of drinking water.

API: Bromhexine HCl 10mg/ml



Pack size: 1L

Mycotect

Indications

Deactivates mycotoxins, restores the intestinal pH to the optimal value and assists in controlling diarrhoea.

Dosages

Chickens: 100ml / 1000 litres of drinking water for 3 to 10 days until mould and fungi infections have ceased.

API: Poly(1-(2-oxo-1-pyrroli-dinyl)ethylene) 3mg/ml



Pack size: 1L

Nutrimax Plus

Indications

Nutrimax Plus contains vitamins and amino acids for optimal livestock growth.

Dosages

Poultry: 1L mix with 1000 to 2000 litres (200 to 400 Gallons) of drinking water.

Sheep, goats, pigs: 5ml / animal

Cattle and horses: 20ml to 30ml / animal

API: Vitamin A 20,000,000 IU/L, Vitamin E 10,000mg/L, Vitamin B1 2,500mg/L, Vitamin B2 1,125mg/L, Vitamin D3 200,000 IU/L, Vitamin B6 500mg/L, Vitamin B12 5mg/L, Vitamin K3 500mg/L, Vitamin B5 2,500mg/L, Biotin 15mg/L, Choline Chloride 50g/L, Lysine 10g/L, Methionine 10g/L, Cystine 6.5g/L, Threonine 5g/L, Leucine 3g/L, Isoleucine 3g/L, Valine 3g/L, Phenylalanine 1.25g/L, Histidine HCl 1g/L, Arginine 3g/L, Aspartame 2g/L, Serine 1.4g/L, Aspartic Acid 2.35g/L, Glutamic Acid 10g/L, Glycine 1.46g/L, Alanine 3.9g/L, Tyrosine 0.7g/L, Proline 1g/L



Pack size: 5L

Quinosol 20%

Indications

For use in turkeys, broilers and pullets being reared as layers for diseases of the respiratory and alimentary tract of bacterial or mycoplasmal origin such as pasteurellosis, mycoplasmosis, colibacillosis, colisepticaemia and salmonellosis.

Dosages

10mg / kg body weight daily.
 Total number of birds x Average body weight in kg x 0.05 =
 Total volume of Quinosol 20% / day.

For the treatment of Salmonellosis and Pasteurellosis:
 5 to 10 days.

For other conditions: 3 to 5 days.

API: Enrofloxacin 200mg/ml



Pack size: 500ml

Taikosol 30%

Indications

Chickens: Prevention and treatment of bacterial infections associated with *Salmonella* spp. and *E. coli* susceptible to florfenicol.

Pigs: Prevention and treatment of bacterial infections associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Mycoplasma pneumoniae* (Mycoplasma) and other respiratory diseases, *Salmonella typhimurium*, and *Streptococcus suis* susceptible to florfenicol.

Dosages

Chickens: 0.067ml / kg body weight for 5 days. This dosage rate is equivalent to 33.34ml / 100L of drinking water for 5 days.

Pigs: 0.34ml / 10kg body weight for 5 days. This dosage rate is equivalent to 33.34ml / 100L of drinking water for 5 days.

API: Florfenicol 300mg/ml



Pack size: 500ml

Thiamisol 200mg/ml

Indications

For the treatment of infectious coryza and respiratory mycoplasma disease caused by *Staphylococcus*, *Streptococcus*, *Corynebacterium*, *Pasteurella*, *Mycobacterium avium* and *Mycoplasma* species in chickens.

Dosages

Chickens (exclude laying hens): 0.5ml to 2.5ml of product / 1L of drinking water.

API: Thiamphenicol 200mg/ml



Pack size: 500ml

Tridiasol 48%

Indications

Pigs: Treatment and control of diseases sensitive to trimethoprim and sulphadiazine, in particular the respiratory disease complex.

Chickens: Treatment and control of diseases sensitive to trimethoprim and sulphadiazine, in particular enteric and respiratory bacterial infections.

Dosages

30mg combined active ingredients / kg body weight daily, given in 2 equal doses.

Chickens: 1.56ml / 25kg body weight daily for up to 7 days.

Pigs: 62.5ml / 1000kg pig weight daily for 5 days.

API: Trimethoprim 80mg/ml, Sulphadiazine 400mg/ml



Pack size: 1L

Trimetosol 40/200mg/ml

Indications

Fattening pigs: Treatment and metaphylaxis of: Post-weaning diarrhoea caused by beta-haemolytic K88-positive, K99-positive or 987P *Escherichia coli* strains susceptible to trimethoprim-sulfamethoxazole. Secondary bacterial infections caused by *Pasteurella multocida*, *Actinobacillus pleuropneumoniae*, *Streptococcus* spp. and *Haemophilus parasuis* susceptible to trimethoprim-sulfamethoxazole.

Broilers: Treatment and metaphylaxis of: Colibacillosis caused by *Escherichia coli* susceptible to trimethoprim-sulfamethoxazole. Coryza caused by *Avibacterium paragallinarum* susceptible to trimethoprim-sulfamethoxazole.

Dosages

Fattening pigs: 5mg trimethoprim and 25mg sulfamethoxazole / kg body weight a day, for 4 to 7 days. This corresponds to 0.5ml of product / 4kg body weight / day.

Broilers: 7.5mg trimethoprim and 37.5mg sulfamethoxazole / kg body weight a day, for 3 days. This corresponds to 0.5ml of product / 2.67kg body weight / day.

API: Trimethoprim 40mg/ml, Sulfamethoxazole 200mg/ml



Pack size: 1L

Trimetosol 80/400mg/ml

Indications

Treatment of bacterial diseases and diarrhoea caused by *E. coli* susceptible to sulfamethoxazole and trimethoprim in piglets.

Dosages

Piglets: 0.125ml of product / 2kg of body weight is administered orally once a day, for 3 to 5 days.

API: Trimethoprim 80mg/ml, Sulfamethoxazole 400mg/ml



Pack size: 500ml, 1L

Vetricox 5mg/ml

Indications

For the treatment of Coccidiosis caused by *E. tenella*, *E. maxima* and *E. acervulina* in broiler chickens.

Dosages

Dilute 0.25ml to 0.5ml (0.05ml to 0.1ml / kg body weight / day) of product / 1L of drinking water and administered for 2 consecutive days.

API: Diclazuril 5mg/ml



Pack size: 200ml, 500ml, 1L

WATER SOLUBLE POWDER

Amprolox 20%

Indications

Poultry: For the treatment of coccidiosis.

Calves: As an aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii*.

Dosages

Poultry:

0.6g Amprolox 20% / litre of fresh drinking water for 3 to 5 days.


In severe outbreaks: 1.2g Amprolox 20% / litre of drinking water. Continue with 0.3g / litre of drinking water for an additional 1 to 2 weeks.

Calves:

5-day treatment protocol: 0.05g Amprolox 20% / kg body weight daily for 5 days.

21-day treatment period: 0.025g Amprolox 20% / kg body weight for 21 days.

API: Amprolium (as HCl) 200mg/g



Pack size: 500g

Aprasure 50%

Indications

Treatment of bacterial enteritis associated with organisms susceptible to apramycin in pigs, colibacillosis and salmonellosis in calves and *E. coli* septicaemia in young chickens.


Dosages

Pigs: 7.5mg to 12.5mg of apramycin / kg body weight daily for 7 days. This may be achieved by adding 2g of Aprasure 50% in 20L of water.

Calves: 2g to 4g Aprasure 50% / 50kg body weight daily for 5 days in drinking water, milk or milk replacer.

Chickens: Mix 500g Aprasure 50% to approximately 500 to 1000 litres (110 to 220 Gallons) of drinking water, for 5 days.

API: Apramycin (as Sulphate) 500mg/g



Pack size: 500g

Aspisure 50%


Indications

For the prevention of heat stress in poultry.

Dosages

Poultry: 0.6g to 1g Aspisure 50% / 10kg body weight 1 to 2 times a day in milk, liquid food or drinking water for 2 to 3 days.

API: Aspirin 500mg/g



Pack size: 500g

Cefalexol 50%

Indications

Effective against the infection of: Respiratory tract, urogenital tract, skin, localised soft tissues and gastrointestinal tract. Broad spectrum antibiotic and effective against the following microorganisms *in vitro*: *Staphylococcus* spp (including penicillin-resistant strains), *Streptococcus* spp., *Corynebacterium* spp., *Pasteurella multocida*, *Escherichia coli* and *Micrococcus* spp.

Dosages

Dilute 200g of product in 1L of drinking water. The recommended dose rate is 1ml to 1.5ml Cefalexol 50% / 10 kg body weight twice daily. Treatment for 5 days is recommended.

The following is intended as a guide:

Dogs: 11kg to 20kg body weight: 1.0ml to 3.0ml twice daily.

6kg to 10kg body weight: 0.5ml to 1.5ml twice daily.

5kg and less body weight: 0.5ml to 1.0ml twice daily.

Cats: Average dose rate: 0.5ml twice daily.

Poultry/Birds:

Prevention: Mix 100g to approximately 400 litres (90 Gallons) of drinking water.

Treatment: Mix 100g to approximately 200 litres (45 Gallons) of drinking water.

Swine, calves, lambs, goats:

Prevention: Mix 1g to 6 litres of drinking water, equivalent to 1g / 45kg to 50kg body weight.

Treatment: Mix 1g to 3 litres of drinking water, equivalent to 1g / 25kg body weight.

API: Cefalexin (as Monohydrate) 500mg/g



Pack Size: 100g, 500g

Cosucin


Indications

Treatment of infections caused by bacteria sensitive to trimethoprim/sulfachloropyridazine, taking into account the ability of the antibiotic to reach the site of infection in effective concentrations.

Dosages

Chickens: 32mg of product / kg of body weight / day (equivalent to 24mg of active substance in total). Particularly in the case of primary and secondary infections by *E. coli*, a higher dose is recommended.

API: Sulfachloropyridazine 625mg/g, Trimethoprim 125mg/g



Pack Size: 100g, 200g, 500g, 1kg

Ctsure 50%

Indications

Chickens: For the treatment and control of colibacillosis secondary to infectious bursal disease, chronic respiratory disease caused by *Mycoplasma gallisepticum* and *Escherichia coli*, *Salmonella enteridis*, *Salmonella typhimurium* and *Pasteurella multocida* infections.

Pigs: For the treatment and control of respiratory diseases associated with *Mycoplasma hyopneumoniae*, *Streptococcus suis* and toxigenic strains of *Pasteurella multocida*. It is also used in the control and treatment of rhinitis due to *Bordetella bronchiseptica* and streptococcal meningitis due to *Streptococcus suis* type II.


Dosages

Chickens: 100g of product should be dissolved in 300L of drinking water and administered daily for up to 5 days. This will provide a daily dosage of 20mg to 50mg chlortetracycline HCl / kg depending on water consumption.

Dosage may be adjusted to up to 60mg / kg depending upon the severity of infection, by careful calculation of the total body weight of the birds and dissolving the requisite amount of CTSURE in the quantity of water consumed within 24 hours.

Pigs: In order to achieve the recommended dosage rate of 20mg chlortetracycline HCl / kg, 100g of product should be dissolved in 250L of water and administered daily for up to 5 days. This will provide medication for 5000kg of pigs.

API: Chlortetracycline HCl 500mg/g



Pack Size: 100g, 200g, 500g, 1kg

Detoxzymes Plus


Indications

Detoxzymes Plus is a multifunctional growth promoting enzyme which eliminates aflatoxins and improves digestion.

Dosages

Poultry: Dissolve 500g of Detoxzymes Plus in 1000 litres to 2000 litres of drinking water. Use daily.

API: Fermentation extracts containing protease and oxidase



Pack Size: 500g, 1kg

Doxsure 50%

Indications

Chickens (broilers): For the treatment of Chronic Respiratory Disease (CRD) caused by *Mycoplasma gallisepticum*.


Fattening pigs: For the treatment of clinical respiratory infection caused by sensitive strains of *Pasteurella multocida*. In case of any infective process, a bacteriological confirmation of the diagnosis is recommended, as well as a sensitivity test of the bacteria causing the process.

Dosages

Chickens (broilers): 40mg of Doxsure 50% Powder (equivalent to 20mg of doxycycline) / kg body weight / day for 3 to 5 days.

Fattening pigs: 20mg of Doxsure 50% Powder (equivalent to 10mg of doxycycline) / kg body weight / day for 5 days.

API: Doxycycline (as HCl) 500mg/g



Pack Size: 500g

D-Stress

Indications


D-Stress is a multivitamin electrolyte product formulated to supply essential vitamins and electrolytes to combat stressful conditions in livestock.

Dosages

Poultry, cattle, sheep and goats:
Water tanks and barrels: 100g in 200L / 44 UK Gallons.

Automatic dosers: Dissolve 375g in 4L of water, set medicator to deliver 40ml of solution per 4L of drinking water.

API: Vitamin A 1,000,000 IU/100g, Vitamin D 200,000 IU/100g, Vitamin E 300mg/100g, Vitamin B6 200mg/100g, Vitamin B12 1mg/100g, Vitamin C 2g/100g, Folic Acid 30mg/100g, Sodium 2.2g/100g, Potassium 0.24g/100g, Magnesium 0.1g/100g, Calcium 0.22g/100g, Chloride 3.3g/100g, Acetate 1.1g/100g, Sulphate 0.4g/100g



Pack Size: 100g

Erytrosure 50%


Indications

Treatment of Chronic Respiratory Disease (CRD) in chickens caused by *Mycoplasma* organisms.

Dosages

33g Erytrosure 50% Powder / 64 litres of drinking water administered continuously for 1 to 5 days, depending on response (approximately 25.5mg erythromycin / kg body weight / day).

API: Erythromycin (as Thiocyanate) 500mg/g



Pack Size: 500g

Lincosure 400mg/g

Indications

Pigs: Treatment and metaphylaxis of enzootic pneumonia caused by *Mycoplasma hyopneumoniae*. The presence of the disease in the group must be established before the product is used.

Chickens: Treatment and metaphylaxis of necrotic enteritis caused by *Clostridium perfringens*. The presence of the disease in the group must be established before the product is used.


Dosages

For use in drinking water.

Pigs: Enzootic pneumonia: 25mg product / kg body weight / day for 21 consecutive days.

Chickens: Necrotic enteritis: 12.5mg product / kg body weight / day for 7 consecutive days.

API: Lincomycin (as HCl) 400mg/g



Pack Size: 100g, 500g

Linco Specsure 222/444mg/g

Indications

Poultry: As an aid in the prevention and treatment of Chronic Respiratory Disease (CRD, Air sacculitis) associated with mycoplasmal and coliform infections of growing chickens.

The presence of the disease in the group must be established before the product is used.


Dosages

Poultry: Add 150g product to 120 litres of drinking water. For automatic medicators set to deliver 7.8ml of stock solution / litre of drinking water, prepare stock solution by dissolving 150g product in 0.9 litres of water.

Prevention: To prevent losses due to CRD, administer at the above dosage for the first 3 to 5 days of life and again for one or two days at about 4 weeks of age. The longer period (5 days) of initial administration is recommended if disease has been a problem in the past on the premises.

Treatment: Begin treatment at the above dosage at the first sign of disease. Administer as the sole source of drinking water for 5 to 7 days. Fresh medicated drinking water should be provided daily.

API: Lincomycin (as HCl) 222mg/g, Spectinomycin (as HCl) 444mg/g



Pack Size: 100g, 500g, 1kg, 25kg

Moxisure 50%

Indications

Chickens, turkeys and ducks: Treatment of infections caused by bacteria susceptible to amoxicillin.

Pigs: For the treatment of Salmonellosis and Pasteurellosis.

Dosages


Chickens: 30mg Moxisure 50% / kg body weight / day. Total treatment period should be 3 days or in severe cases for 5 days.

Ducks: 40mg Moxisure 50% / kg body weight / day for 3 days.
Turkeys: 30mg to 40mg Moxisure 50% / kg body weight / day for 3 days or in severe cases for 5 days.

Pigs: For administration via drinking water: 40mg Moxisure 50% / kg body weight / day, administered at 24 hour intervals for up to 5 days.

For administration in liquid feed: 40mg Moxisure 50% / kg body weight / day for up to 5 days.

API: Amoxicillin Trihydrate 500mg/g



Pack Size: 500g, 1kg

Neocin 49%

Indications


Treatment of gastrointestinal infections caused by bacteria sensitive to neomycin in piglets and poultry.

Dosages

Piglets: 2.04g to 3.06g Neocin 49% / 100kg body weight every 12 hours for 3 to 5 consecutive days.

Poultry: 4.08g to 6.12g Neocin 49% / 10 litres of drinking water for 3 to 5 consecutive days.

API: Neomycin (as Sulphate) 490mg/g



Pack Size: 500g

Oxysure 50%

Indications

Oxytetracycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria like *Bordetella*, *Bacillus*, *Corynebacterium*, *Campylobacter*, *E. coli*, *Haemophilus*, *Pasteurella*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. and *Mycoplasma*, *Rickettsia* and *Chlamydia* spp.

The mode of action of oxytetracycline is based on inhibition of bacterial protein synthesis. Oxytetracycline is mainly excreted in urine and to a lesser degree in bile and in lactating animals in milk.


Dosages

Chickens and turkeys: 2g of Oxysure 50% / 20L of drinking water for 3 to 5 days.

For use in feed: For control of bacterial diseases, add 200g of Oxysure 50% / metric tonne of feed.

For treatment of bacterial diseases, add 0.4kg to 0.8kg of Oxysure 50% / metric tonne of feed.

API: Oxytetracycline HCl 500mg/g



Pack Size: 500g

Sulfacox 50%


Indications

Chickens: Treatment for coccidiosis.

Dosages

0.03% of Sulfacox 50% in drinking water. Administer in drinking water for 3 days as sole source of drinking water and sulphonamide medication.

API: Sulfachloropyrazine Sodium Monohydrate 500mg/g



Pack Size: 108g

Taipan

Indications

Aids growth and digestion, stimulates appetite, supports nutrient absorption, and helps immunity. Prevention of diseases due to vitamin deficiencies, irregular digestion or digestion failure.

Dosages

Poultry, swine, cattle, sheep and goats:

1g in 1 litre of drinking water.

1kg to 2kg / tonne of feed.

Fish and shrimp: 1kg to 2kg / tonne of feed.

API: Vitamin A 3,000,000 IU/kg, Vitamin D3 600,000 IU/kg, Vitamin E 1,000mg/kg, Vitamin B1 1,000mg/kg, Vitamin B2 1,000mg/kg, Vitamin B6 500mg/kg, Nicotinamide 10,000mg/kg, Calcium D-Pantothenate 2,000mg/kg, Vitamin B12 1,000mcg/kg, Folic Acid 100mg/kg, Vitamin C 30,000mg/kg, L-Lysine 10,000mg/kg, Threonine 500mg/kg, DL-Methionine 10,000mg/kg, Biotin 1,000mcg/kg, Noritase 60,000mg/kg



Pack Size: 500g, 1kg

Tylosure 500mg/g

Indications

Calves: For the treatment and metaphylaxis of pneumonia due to *Mycoplasma* spp.

Pigs: Treatment and prevention of enzootic pneumonia and ileitis.

Chickens: Treatment and prevention of CRD and necrotic enteritis.

Turkeys: Treatment and prevention of infectious sinusitis.

Dosages

Calves: Pneumonia: 2g to 4g product / 100kg body weight 2 times daily for 7 to 14 days.

Pigs: Treatment of enzootic pneumonia: 40mg of Tylosure 50% / kg body weight / day for 10 days.


Treatment of ileitis or PIA: 10mg to 20mg of Tylosure 50% / kg body weight / day for 7 days.

Chickens: Treatment of chronic respiratory disease: 150mg to 200mg of Tylosure 50% / kg body weight / day for 3 to 5 days.

Treatment of necrotic enteritis: 40mg of Tylosure 50% / kg body weight / day for 3 days.

Turkeys: 150mg to 200mg of Tylosure 50% / kg body weight / day for 3 to 5 days.

API: Tylosin (as Tartrate) 500mg/g



Pack Size: 500g

Tylvalosure 625mg/g

Indications

Chickens: Treatment and metaphylaxis of respiratory infections caused by *Mycoplasma gallisepticum*. The presence of the disease in the flock should be established before metaphylactic treatment. As an aid in reducing the development of clinical signs and mortality from respiratory disease in flocks, where infection in ovum with *Mycoplasma gallisepticum* is likely because the disease is known to exist in the parent generation.

Turkeys: Treatment of respiratory disease associated with tylvalosin sensitive strains of *Ornithobacterium rhinotracheale*.


Dosages

Chickens: For treatment: 25mg tylvalosin / kg body weight / day in drinking water for 3 consecutive days.

For prevention: 25mg tylvalosin / kg body weight / day in drinking water for 3 consecutive days at 1 day old. This is followed by a second treatment with 25mg tylvalosin / kg body weight / day in drinking water for 3 consecutive days at the period risk, i.e. at times of management stress such as administration of vaccines (typically when birds are 2 to 3 weeks old).

Turkeys: 25mg tylvalosin / kg body weight / day in drinking water for 5 consecutive day.

API: Tylvalosin (as Tartrate) 625mg/g



Pack Size: 100g, 500g

VAM Plus

Indications


VAM Plus contains 13 vitamins, 18 amino acid and 9 minerals for optimal livestock growth.

Dosages

Poultry, cattle, sheep and goats:

Water tanks and barrels: 100g in 50 to 100 UK Gallons of water (230L to 454L).

API: Vitamin A 13,500,000 IU/kg, Vitamin D 1,250,000 IU/kg, Vitamin E 7,500 IU/kg, Vitamin B1 1.3g/kg, Vitamin B2 2.5g/kg, Vitamin B6 3g/kg, Vitamin B12 12.5mg/kg, Vitamin C 7.5g/kg, Vitamin H 5mg/kg, Calcium Pantothenate 7.5g/kg, Folic Acid 250mg/kg, Nicotinamide 30g/kg, Inositol 0.7g/kg, Lysine 10.9g/kg, Methionine 1.0g/kg, Cystine 0.3g/kg, Threonine 5.6g/kg, Tryptophan 2.0g/kg, Leucine 10g/kg, Isoleucine 6.6g/kg, Valine 7.0g/kg, Phenylalanine 5.5g/kg, Histidine 2.8g/kg, Arginine 2.9g/kg, Aspartic Acid 12.6g/kg, Serine 5.6g/kg, Glutamic Acid 17.6g/kg, Glycine 6.3g/kg, Alanine 9.1g/kg, Tyrosine 4.6g/kg, Proline 6.3g/kg, Calcium 200mg/kg, Phosphorus 3.72mg/kg, Sodium 684mg/kg, Magnesium 436mg/kg, Potassium 4.8mg/kg, Copper 200mcg/kg, Zinc 12.2mg/kg, Iron 16mg/kg, Manganese 2.5mg/kg



Pack Size: 100g

Vibrasure 500mg/g

Indications

Treatment of the following specified infectious diseases of the respiratory tract and the alimentary tract caused by microorganisms susceptible to doxycycline.

Cattle (Pre-ruminant calves):

Bronchopneumonia and pleuropneumonia caused by *Pasteurella* spp., *Streptococcus* spp., *Trueperella pyogenes*, *Histophilus somni* and *Mycoplasma* spp.

Pigs:

Atrophic rhinitis caused by *Pasteurella multocida* and *Bordetella bronchiseptica*; Bronchopneumonia caused by *Pasteurella multocida*, *Streptococcus suis* and *Mycoplasma hyorhinis*; Pleuropneumonia caused by *Actinobacillus pleuropneumoniae*.

Chickens (broilers, breeders, replacement pullets):

Infections of the respiratory tract caused by *Mycoplasma* spp., *Escherichia coli*, *Haemophilus paragallinarum* and *Bordetella avium*; Enteritis caused by *Clostridium perfringens* and *Clostridium colinum*.

Dosages


To be administered orally through the milk-replacer and/or the drinking water.

Cattle (pre-ruminant calves): For use in milk replacer. 20mg of product / kg body weight / day, divided over 2 administrations, for 3 to 5 consecutive days.

Pigs: For use in drinking water. 20mg of product / kg body weight / day, for 3 to 5 consecutive days.

Chickens (broilers, breeders, replacement pullets): For use in drinking water. 50mg of product / kg body weight / day, for 3 to 5 consecutive days.

API: Doxycycline Hyclate 500g/kg



Pack Size: 100g, 500g

Zeolytes


Indications

Specially formulated to provide rapid correction of dehydration, electrolyte loss and metabolic acidosis particularly when these disturbances arise as a consequence of diarrhoea or during heat stress in swine.

Dosages

Poultry: 500g to be dissolved in 450 to 900 litres of drinking water. Use for 3 to 5 days.

API: Sodium 0.264g/g, Potassium 0.08g/g, Magnesium 0.8mg/g, Bicarbonate 0.41g/g, Chloride 0.238g/g, Acetate 2.2mg/g



Pack Size: 500g, 1kg

ORAL PUMPS

Enroxon 5mg/ml

Indications

For the treatment of diseases of the respiratory and alimentary tracts of bacterial or mycoplasmal origin (e.g. pasteurellosis, mycoplasmosis, colibacillosis, colisepticaemia and salmonellosis), and multifactorial diseases such as atrophic rhinitis and enzootic pneumonia, where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the drug of choice. and enzootic pneumonia, where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the drug of choice.

Dosages

1.5mg to 5mg enrofloxacin / kg body weight daily.
For piglets up to 3kg body weight: 1ml once daily for 3 to 5 days.
For piglets 3kg up to 10kg body weight: 3ml once daily for 3 to 5 days. 1 pump stroke delivers 1ml of product.

API: Enrofloxacin 5mg/ml



Pack Size: 200ml

Gentacure 0.5%

Indications

Control and treatment of colibacillosis in neonatal pigs (1 to 3 days old) caused by *E. coli* sensitive to gentamicin.

Dosages

One full pump (1ml Gentacure 0.5%) / pig one time at the first sign of disease.

API: Gentamicin (as Sulphate) 5mg/ml



Pack Size: 200ml

Oradex 10% + Oralgen 6.4%

Indications

Oradex 10%:
Prevention of iron deficiency anaemia in piglets.

Oralgen 6.4%:
Treatment of *E. coli* bacterial diarrhoea in piglets.

Dosages

Dilute 5ml Oralgen 6.4% in 200ml of Oradex 10%. Shake well and administer 2ml / piglet within 24 hours of farrowing. One pump delivers 1ml of solution.

**API: Oradex 10%: Iron (as Dextran) 100mg/ml,
Oralgen 6.4%: Gentamicin 64000 IU/ml**



Pack Sizes: Oradex 200ml
Oralgen 5ml

Pigicox 5%

Indications

Pigs: Prevention of clinical signs of coccidiosis in neonatal piglets on farms with a confirmed history of coccidiosis caused by *Cystoisospora suis*.

Dosages

Toltrazuril 20mg / kg body weight, corresponding to 1ml Pigicox 5% (1 pump) / 2.5 kg body weight for 3 to 5 days.

API: Toltrazuril 50mg/ml



Pack Size: 200ml

FEED ADDITIVES

Amprofeed 250g/kg Powder

Indications

Turkeys: As an aid in the prevention of coccidiosis.

Broiler chickens: As an aid in the prevention of deaths from coccidiosis.

Replacement chickens: As an aid in the prevention of death from coccidiosis in chickens intended as conventional floor managed layers. As an aid in the prevention of deaths from coccidiosis in chickens intended as caged layers.

Calves: As an aid in the prevention and treatment of coccidiosis caused by *Eimeria bovis* and *E. zuernii*.

Dosages

Turkeys: 500g of product / tonne of feed as sole ration from one day of age up to and including 16 weeks of age.

Broiler chickens: 500g or 1kg of product / tonne of feed as sole ration from one day of age until birds are slaughtered.

Replacement chickens:

Chickens intended as conventional floor managed layers:

Heavy exposure to coccidiosis:

Age 0-5 weeks: 500g of product / tonne of feed

Age 5-9 weeks: 320g to 500g of product / tonne of feed

Age 9-14 weeks: 320g of product / tonne of feed

Light to moderate exposure to coccidiosis:

Age 0-5 weeks: 320g of product / tonne of feed

Age 5-9 weeks: 240g to 320g of product / tonne of feed

Age 9-14 weeks: 240g of product / tonne of feed

Chickens intended as caged layers: 500g or 1kg of product / tonne of feed as sole ration from one day of age until birds are placed in cages.

Calves:

Prevention: 250mg of amprolium / 50kg of body weight (5mg / kg) / day for 21 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

Treatment: 500mg of amprolium / 50kg of body weight (10mg / kg) / day for 5 days. The proper amount of amprolium medicated supplement is mixed with the amount of ration usually consumed in one day.

API: Amprolium 250g/kg



Pack Sizes: 25kg

Candistan 4.4% Powder

Indications

Control and treatment of crop mycosis and mycotic diarrhoea (*Candida albicans*).

Dosages

Control of crop of mycosis and mycotic diarrhoea (*Candida albicans*):
1.136kg Candistan 4.4% (equivalent to 140 MIU or 50g of nystatin) / tonne of feed.

Treatment of crop mycosis and mycotic diarrhoea (*Candida albicans*):
2.273kg Candistan 4.4% (equivalent to 280 MIU or 100g of nystatin) / tonne of feed for 7 to 10 days.

API: Nystatin 123.2 MIU/kg



Pack Sizes: 25kg

Chick Acid

Indications

For the prevention of indigestion and undigested feed which can lead to diarrhoea and mortality.
For the prevention of microbial overload in the digestive tract.
To prevent the contamination of feed by Salmonella, yeast and mould.

Dosages

Poultry: 1kg to 2kg / tonne of feed.

API: Inorganic acids



Pack Sizes: 25kg

Clopifeed 250g/kg Powder

Indications

Chickens (broilers and layer replacements): Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*.

Turkeys: Aid in the prevention of leucocytozoonosis caused by *Leucocytozoon smithi*.

Dosages

Chickens (broilers and layer replacements): 113.5g clopidol / tonne (907kg) of feed. Do not feed to chickens over 16 weeks of age. If intended for use as caged layers: 227g clopidol / tonne (907kg) of feed. Feed up to 16 weeks of age continuously as the sole ration.

Turkeys: For turkeys grown for meat purposes only: Administer 113.5g or 227g (0.0125 or 0.025 percent) clopidol / tonne (907kg) of feed continuously as the sole ration depending upon management practices, degree of exposure and amount of feed eaten.

API: Clopidol 250g/kg



Pack Sizes: 25kg

Decofeed 60g/kg Powder

Indications

Sheep (young): For the prevention of coccidiosis caused by *Eimeria ovinoidalis*, *E. parva*, *E. bakuensis* and *E. crandallis*.

Chickens (broilers): For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. mivati*, *E. acervulina*, *E. maxima* and *E. brunetti*.

Goats (young): For the prevention of coccidiosis caused by *Eimeria christensenii* and *E. ninakohlyakimovae*.

Cattle: For the prevention of coccidiosis in cattle (including ruminating and non-ruminating calves and veal calves) caused by *Eimeria bovis* and *E. zuerni*.

Dosages

Sheep and goats (young): 0.5mg decoquinate / kg body weight / day. Administer as a top dress supplement or mix into daily ration: 90.9g to 535.7g decoquinate / tonne (907kg) of feed.

Administer in feed type C feed or milk replacer: 12.9g to 90.8g decoquinate / tonne (907kg) of feed. Feed for at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard. Bentonite should not be used in decoquinate feeds.

Chickens (broilers): 27.2g decoquinate / tonne (907kg) of feed. Bentonite should not be used in decoquinate feeds.

Cattle: 27.2g decoquinate / tonne (907kg) of feed.

API: Decoquinate 60g/kg



Pack Size: 25kg

Detoxzymes Pro

Indications

Detoxzymes Pro is a perfectly-selected protein for the performance of livestock.

Dosages

To be used daily.

Swine:

Pre-starters and starters: 0.5kg to 1kg / tonne of feed.

Growers and finishers: 0.5kg / tonne of feed.

Sows: 0.5kg / tonne of feed.

API: *Saccharomyces cerevisiae* fermentation extract containing polysaccharides and proteins in the form of peptides.



Pack Size: 25kg

Dotfeed 250g/kg Powder

Indications

An aid in the prevention of coccidiosis of broilers, replacement pullets and turkeys.

Dosages

Broilers and replacement pullets: Thoroughly mix 500g of product / tonne of feed to produce a concentration of 125g dinitolmide / tonne in the finished feed.

Turkey growers: Thoroughly mix 760g of product / tonne of feed to produce a concentration of 190g dinitolmide / tonne in the finished feed.

API: Dinitolmide 250g/kg



Pack Size: 25kg

Drisezymes Powder

Indications

Improves the digestive system, reduces diarrhoea outbreaks, increases absorption of nutrients, improve carcass quality, improves appetite and long-term health, reduces mortality, improves feed conversion and growth rate in fish and livestock.

Dosages

Piglets: 1.0kg / tonne of feed

Pigs: 0.5kg / tonne of feed

Poultry: 0.5kg / tonne of feed

Fish: 0.5kg / tonne of feed

API: Alpha-amylase 6000 KNU/kg, Protease 22.5 AU/kg



Pack Size: 25kg

Duracin 1% Powder

Indications

For the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. maxima* and *E. mivati*.

Dosages

454g to 545g of Duracin 1% Powder which is equivalent to 4.54g to 5.45g (5ppm to 6ppm) maduramicin ammonium / tonne of feed / day for 5 consecutive days orally in feed.

API: Maduramicin Ammonium 10g/kg



Pack Size: 25kg

Em-Micofeed 10% Granules

Indications

Treatment of pneumonia caused by *Actinobacillus pleuropneumoniae*, *Mycoplasma hyopneumoniae*, *Pasteurella multocida* sensitive to tilmicosin in weaned fattening pigs.

Dosages

8mg to 16mg of Em-Micofeed 10% / kg body weight / day, corresponding to 2kg to 4kg / tonne of feed for 15 days.

API: Tilmicosin (as Phosphate) 100g/kg



Pack Size: 25kg

Em-Taikomix 10% Granules

Indications

Treatment and prevention of swine respiratory disease in infected herds due to *Pasteurella multocida* susceptible to florfenicol.

Dosages

100mg Em-Taikomix 10% / day for 5 consecutive days. For a daily feed intake of 50g feed / kg body weight, this dosage corresponds to a rate of incorporation of 2kg of premix / tonne of feed.

API: Florfenicol 100g/kg



Pack Size: 10kg, 25kg

Florosure 50% Powder

Indications

For the treatment and prevention of swine respiratory disease in infected herds due to *Pasteurella multocida* susceptible to florfenicol.

Dosages

10mg of florfenicol / kg body weight (equivalent to 20mg Florosure 50%) / day administered for 5 consecutive days.

For a daily feed intake of 50g / kg body weight, this dosage corresponds to a rate of incorporation of 0.4kg of Florosure 50% / tonne of feed, i.e. 200 ppm of florfenicol.

API: Florfenicol 500g/kg



Pack Size: 10kg, 25kg

Kollitides Pro

Indications

Kollitides Pro is a fermentation extract containing high amount of nutritional components that are essential to the growth of normal flora in the gut of animal.

Dosages

Broilers, layers, ducks and pigs: 1kg to 2kg / tonne of feed.

API: *Saccharomyces cerevisiae* fermentation extract containing polysaccharides and proteins in the form of peptides.



Pack Size: 10kg, 25kg

Linco Actinmix Powder

Indications

Pigs: Control and treatment of swine dysentery, enteritis, complicated or mixed enteric infections, and ileitis.
An aid in the control of mycoplasmal (enzootic) pneumonia.
Treatment of Mastitis, Metritis, Agalactia (MMA) syndrome of bacterial origin in sows.

Dosages

Enteric disease:

Control: 1kg to 2kg of Linco Actinmix / tonne of feed over period of risk.

Treatment: 2kg / tonne of feed for 3 weeks or until the clinical signs disappear.

Control of mycoplasmal pneumonia: 1kg to 2kg of Linco Actinmix / tonne of feed, feed daily throughout the period of risk.

Treatment of MMA: 1kg to 2kg of Linco Actinmix / tonne of feed for 5 to 10 days prior to farrowing and 2 to 3 weeks post-farrowing.

API: Lincomycin (as HCl) 22g/kg, Spectinomycin (as HCl) 22g/kg



Pack Size: 25kg

Lincofeed 4.4% Powder

Indications

For treatment and control of swine dysentery and mycoplasmal pneumonia.

Dosages

Control of swine dysentery: Add 1kg to 1000kg of feed as the sole ration throughout the period of risk.

Treatment of swine dysentery: Add 2.5kg to 1000kg of feed as the sole ration for three weeks or until clinical signs of disease (watery, mucoid or bloody stools) disappear.

Treatment and control of mycoplasmal pneumonia: Add 5kg to 1000kg of feed as the sole ration for three weeks or until clinical signs of disease disappear.

API: Lincomycin (as HCl) 44g/kg



Pack Size: 10kg, 25kg

Lipotide

Indications

A new antimicrobial peptide to combat Gram-negative bacteria.

Dosages

To be used daily.

Broiler (starter): 100g to 150g / tonne feed.

Broiler (grower/finisher): 150g to 200g / tonne feed.

API: Fermentation extract containing polysaccharides and proteins in the form of peptides.



Pack Size: 20kg

Microtide

Indications

A new antimicrobial peptide to combat Gram-negative bacteria.

Dosages

To be used daily.

Broiler (starter): 200g to 250g / tonne feed.

Broiler (grower/finisher): 150g to 250g / tonne feed.

API: Fermentation extract containing polysaccharides and proteins in the form of peptides



Pack Size: 20kg

Milk Acid

Indications

For the prevention of indigestion and undigested feed which can lead to fermentation, diarrhoea and mortality.
For the prevention of insufficient chlorhydropeptic secretions and enzymatic secretions of the digestive tract as well as the prevention of microbial overload in the digestive tract.

Dosages

Prestarters, starters, growers and sows: 1kg to 2kg Milk Acid / tonne of feed.

API: Orthophosphoric Acid, Malic Acid, Citric Acid, Lactic Acid, Tartaric Acid



Pack Size: 25kg

Moxifeed 500g/kg Granules

Indications

Broiler chickens: For the prevention of respiratory infections caused by *E. coli* susceptible to amoxicillin.

Dosages

Broiler chickens: 20mg of amoxicillin equivalent to 23mg amoxicillin trihydrate (46mg of products) / kg body weight / day for 5 consecutive days in feed.

API: Amoxicillin Trihydrate 500g/kg



Pack Size: 10kg, 25kg

Neofeed 490,000 IU/g Powder

Indications

For treatment and prevention of gastrointestinal infections due to *Escherichia coli* in pigs, chickens, turkeys and ducks.

Dosages

20,000 IU neomycin / kg body weight / day, orally, for 10 days.

Incorporation rate

The amount of medicated feed consumed by animals depends on their physiological and clinical status. In order to comply with the recommended doses of neomycin and take into account the actual dietary intake, the incorporation rate may be increased or possibly decreased to the minimum incorporation limit of 200 IU neomycin / g of feed. This may lead to levels of neomycin in the medicated feed different from those mentioned below for information purposes.

Pigs: For daily feed intake of 50g of feed / kg body weight, this dosage corresponds to 400 IU neomycin / gram of feed.

Chickens, turkeys and ducks: For daily feed intake of 100g of feed / kg body weight, this dosage corresponds to 200 IU neomycin / gram of feed.

API: Neomycin Sulphate 490,000 IU/g



Pack Size: 10kg, 25kg

Nicoxin 25% Powder

Indications

Aid in the prevention of coccidiosis in chickens caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. necatrix* and *E. tenella*.

Dosages

Prevention of coccidiosis: 500g / tonne of feed.

API: Nicarbazin 250g/kg



Pack Size: 10kg, 25kg

Quinofeed 200g/kg Granules

Indications

Pigs: Treatment of diarrhoea, enterotoxigenesis, edema and colibacillosis caused by *E. coli* susceptible to enrofloxacin. Treatment of respiratory disease caused by *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and Pleural pneumonia (*Actinobacillus pleuropneumoniae*) susceptible to enrofloxacin.

Dosages

Pigs:

Diarrhoea, enterotoxigenesis, edema and colibacillosis: 0.25kg of product / tonne of feed / day which is equivalent to 50ppm of enrofloxacin uniformly mixed for 5 days. May be given up to 10 days depending on symptoms.

Respiratory disease: 0.75kg of product / tonne of feed / day which is equivalent to 150ppm of enrofloxacin uniformly mixed for 7 days. May be given up to 10 days depending on symptoms.

API: Enrofloxacin 200g/kg



Pack Size: 10kg, 25kg

Robenxin 6.6% Powder

Indications

As an aid in the prevention of coccidiosis caused by *E. mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima* and *E. necatrix* in broiler chickens.

Dosages

Broiler chickens and turkeys: Mix 500g of Robenxin 6.6% / tonne of complete feed.

API: Robenidine HCl 66g/kg



Pack Size: 25kg

Taipan FA

Indications

Aids growth and digestion, stimulates appetite, supports nutrient absorption, and helps immunity. Prevention of diseases due to vitamin deficiencies, irregular digestion or digestion failure.

Dosages

Poultry, swine, cattle, sheep, goats, fish and shrimp:
500g to 2kg / tonne of feed.

API: Vitamin A 3,000,000 IU/kg, Vitamin D3 600,000 IU/kg, Vitamin E 1,000mg/kg, Vitamin B1 1,000mg/kg, Vitamin B2 1,000mg/kg, Vitamin B6 500mg/kg, Nicotinamide 10,000mg/kg, Calcium D-Pantothenate 2,000mg/kg, Vitamin B12 1,000mcg/kg, Folic Acid 100mg/kg, Vitamin C 30,000mg/kg, L-Lysine 10,000mg/kg, Threonine 500mg/kg, DL-Methionine 10,000mg/kg, Biotin 1,000mcg/kg, Noritase 60,000mg/kg



Pack Size: 25kg

Trimetomix 40% Powder

Indications

Pigs: Treatment of diseases caused by bacteria sensitive to the combination sulphadiazine and trimethoprim, atrophic rhinitis and reduces mortality due to Streptococcal meningitis.

Non-laying chickens: Treatment of diseases caused by bacteria sensitive to the combination sulphadiazine and trimethoprim, such as *E. coli*, *Salmonella* spp. and *Pasteurella multocida*.

Dosages

Pigs: 0.75kg to 1.13kg Trimetomix 40% / tonne of feed for 5 consecutive days.

Non-laying chickens: 0.75kg Trimetomix 40% / tonne of feed for 10 consecutive days.

API: Trimethoprim 66.7g/kg, Sulphadiazine 333.3g/kg



Pack Size: 25kg

Trimetosol 80/400g/kg Powder

Indications

Treatment of bacterial diseases and diarrhoea caused by *E. coli* susceptible to trimethoprim and sulfamethoxazole in piglets and chickens.

Dosages

The following dose is mixed with feed and administered orally once a day for 5 to 7 days. 551g of product / tonne of feed.

API: Trimethoprim 80g/kg, Sulfamethoxazole 400g/kg



Pack Size: 25kg

Zuricox 0.5% Powder

Indications

Broiler chickens. Aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mitis*, *E. necatrix* and *E. tenella*.

Growing turkeys. As an aid in the prevention of coccidiosis caused by *Eimeria adenoeides*, *E. meleagrimitis* and *E. gallopavonis*.

Dosages

Broiler chickens: 200g Zuricox 0.5% / tonne of feed (equivalent to 1 part per million diclazuril) continuously.

Growing turkeys: 200g Zuricox 0.5% / tonne of feed (equivalent to 1 part per million diclazuril) continuously.

API: Diclazuril 5g/kg



Pack Size: 25kg

DISINFECTANTS

Germecide

Indications

General use: For the elimination of disease causing organisms by application to building and equipment.

Specific use: For the routine disinfection of premises and equipment to control pathogen causing enteritis, septicaemia and respiratory diseases in farm animals.

Dosages

Basic Use (Terminal & Routine Disinfection):
To be diluted with water.

General dosage: 1 litre of Germecide in 200 litres of water to cover 600 square metre

Dipping/Foot bath: No specific disease – 1 litre of Germecide diluted in 200 litres of water

Terminal disinfection: 2.5 litres of Germecide in 2.5 litres of water. Use this solution of 5 litres for 1000 cubic metre

API: Glutaraldehyde 62.5g/L, Dimethyl Coco Benzyl Ammonium Chloride 50g/L, Didecyl Dimethyl Ammonium Chloride 37.5g/L, Octyl Decyldimethyl Ammonium Chloride 37.5g/L, Pine Oil 10g/L



Pack Size: 5 Litre

Iodophor

Indications

General elimination of disease-causing organisms by application to building and equipment.
For the routine disinfection of premises and equipment to control pathogens causing enteritis, septicaemia and respiratory diseases in farm animals.

Dosages

To be diluted with water.

For cleaning and disinfection of livestock houses and equipment, 1:500 (200ml to 100L).

For disinfection against foot and mouth disease virus, 1:250 (400ml to 100L).

For fogging and spraying poultry houses and equipment against NCD, CRD and infectious coryza, 1:350 (300ml to 100L).

For foot baths and vehicle dips, 1:180 (550ml to 100L).

API: Iodine 30g/L



Pack Size: 5 Litre

Klenox

Indications

To mainly be used as a cleaner and disinfectant of water systems. Removes organic biofilm in the whole drinking system. Kills bacteria, yeasts and fungi in the water.

To be used during empty periods. Klenox effectively decontaminates and disinfects by means of oxidation and is safe to use with minimal risk of damage to equipment. Klenox converts to oxygen and water in the water system, making it a safe option to use for the environment.

Dosages

Usage during operational periods: 50ml / 1000L of water on the first day of use. 100ml / 1000L of water to be used daily after.

Usage during empty periods: 1L to 100L to 250L of water. Leave in water system for 6 hours to 24 hours.

API: Hydrogen Peroxide 50%



Pack Size: 5 Litre

Microcide

Indications

General elimination of disease-causing organisms by application to building and equipment.

For the routine disinfection of premises and equipment to control pathogens causing enteritis, septicaemia and respiratory diseases in farm animals.

Dosages

For routine disinfection in piggeries or poultry units, dilute at a rate of 1:150 to 1:400 and apply 300ml / square meter.

To control major pathogen causing septicaemia, enteritis, respiratory diseases and canine parvovirus, 1:50.

Disinfection of drinking water, 1:3000.

API: Glutaraldehyde 150g/L, Dimethyl Coco Benzyl Ammonium Chloride 100g/L, Pine Oil 50g/L



Pack Size: 5 Litre

Viruxéal

Indications

General use: For use in cleaning and disinfecting associated with livestock facilities.

Dosages

General Fogging: Prepare Viruxéal at dilution of 1% to 2% (1:100 to 1:50). Apply at an applicable rate of 1L to 120 sq. ft. of floor space.

Broiler/Breeder Houses: Spray floors and walls with 1% Viruxéal solution. Thoroughly wash waters and feeders with 1% Viruxéal solution. After contact for 10 minutes, rinse with water.

Processing Plants: Spray with 1% Viruxéal solution to disinfect and clean walls, ceilings and floors.

Swine and Cattle Houses: For most bacteria and viruses, including and not limited to: *Actinobacillus pleuropneumoniae*, *Clostridium perfringens*, foot and mouth disease, ASFV, PRRS, PCV2, spray with 2% Viruxéal solution.

Directions for use: To achieve the appropriate use-diluted Viruxéal concentration, fill container with the volume of water and quantity of powder as per following table. Solutions are stable for 7 days. Do not soak metal objects in Viruxéal for long periods – 10 minutes is maximum necessary contact time. 1L solution is sufficient to treat 33 sq. ft.

Quantity of Water	Dilution Rate			
	1:50	1:100	1:200	Continuous Water Disinfection 1:1000
100L	2kg	1kg	500g	100g
250L	5kg	2.5kg	1.25kg	250g
500L	10kg	5kg	2.5kg	500g

API: Pentapotassium Peroxymonosulphate 21.41%, Sodium Chloride 1.50%



Pack Size: 2kg, 5kg

Viruxeal Aqua

Indications

Disinfecting pond water.

Dosages

For shrimp:

Before stocking: 0.6kg / 1,000m³. Post larvae can be stocked after 1 day.

During culture: 0.5kg to 1kg / 1,000m³.

Periodic use every 10 to 15 days is recommended.

Usage: For shrimp: Dissolve 1kg Viruxeal Aqua in 20L of fresh water then spread evenly on the pond at any time.

Equipment disinfection: Dilute 100g / 10L of water to dip equipment (nets, rackets) within 2 to 3 hours.

API: Potassium Monopersulphate Triple Salt 49.8%



Pack Size: 2kg, 5kg

Contact

Tel: (603) 6156 8708

Fax: (603) 6156 8707

Email: info@rangepharma.com

Address: No. 1, Jalan TSB 11, Taman Industri Sg. Buloh,
47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia.

